




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 :		A1	(11) International Publication Number: WO 00/26203
C07D 277/48, 417/12, 417/14, 471/10, 491/10, A61K 31/426, 31/427, C07D 471/04 // (C07D 471/10, 235:00, 221:00) (C07D 491/10, 317:00, 221:00) (C07D 471/04, 235:00, 221:00)			(43) International Publication Date: 11 May 2000 (11.05.00)
(21) International Application Number: PCT/EP99/08307		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 27 October 1999 (27.10.99)		Published With international search report.	
(30) Priority Data: 9823873.6 30 October 1998 (30.10.98) GB <i>30 Apr 01 / 30 m/s</i>			
(71) Applicant (for all designated States except US): PHARMACIA & UPJOHN S.P.A. [IT/IT]; Via Robert Koch, 1.2, I-20152 Milano (IT).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): <del>DEVARELLO, Paolo</del> [IT/IT]; Piazza San Pietro in Ciel d'Oro, 7/A, I-27100 Pavia (IT), <del>AMICI, Raffaella</del> [IT/IT]; Via N. Rocca, 11, I-29100 Piacenza (IT), <del>TRAQUANDI, Gabriela</del> [IT/IT]; Via F. Cilea, 106, I-20151 Milano (IT), <del>VILLA, Manuela</del> [IT/IT]; Via San Bernardino, 12, I-22040 Lurago d'Erba (IT), <del>VULPETTI, Anna</del> [IT/IT]; Via Volturro Portici/2 80, I-20047 Brugherio (IT), <del>ISACCHI, Antonella</del> [IT/IT]; Via Montecatini, 14, I-20144 Milano (IT).			

(54) Title: 2-UREIDO-THIAZOLE DERIVATIVES, PROCESS FOR THEIR PREPARATION, AND THEIR USE AS ANTITUMOR AGENTS



(I)

## (57) Abstract

Compounds which are 2-ureido-1,3-thiazole derivatives of formula (I) wherein R is a halogen atom, a nitro group, an optionally substituted amino group or it is a group, optionally further substituted, selected from: i) straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl; ii) C<sub>3</sub>-C<sub>6</sub> substituted cycloalkyl; iii) aryl or arylalkyl with from 1 to 6 carbon atoms within the straight or branched alkyl chain; R<sub>1</sub> is an optionally substituted group selected from: i) straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl; ii) 3 to 6 membered carbocycle or 5 to 7 membered heterocycle ring; iii) aryl or arylcarbonyl; iv) arylalkyl with from 1 to 6 carbon atoms within the straight or branched alkyl chain; R<sub>2</sub> is hydrogen, a straight or branched C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>2</sub>-C<sub>4</sub> alkenyl or alkynyl group; or, taken together with the nitrogen atom to which they are bonded, R<sub>1</sub> and R<sub>2</sub> form a 9 to 11 membered spiro-heterocyclic compound; or a pharmaceutically acceptable salt thereof; are useful for treating cell proliferative disorders associated with an altered cell dependent kinase activity.